

## INTRODUCTION


- Globally, up to 95% of adult systemic mastocytosis (SM) cases are associated with somatic gain-of-function point mutations in the *KIT* gene, mainly *KIT* D816V mutations<sup>1-3</sup>
- Little is known about the molecular profile characterization of the *KIT* gene in Europe and whether clinicians are accurately diagnosing SM
- Little is also known about the co-occurrence of somatic variants in other genes such as *SRSF2*, *ASXL1*, *TET2*, and *RUNX1*
- Accurate diagnosis and understanding of co-mutations in SM can impact prognosis and treatment strategy<sup>4,5</sup>

## AIMS

- We explored the landscape of next-generation sequencing (NGS) testing practices for *KIT* mutations in clinical practice in France, Italy, Germany, and Spain, to determine the potential number of individuals that could be impacted by *KIT*-targeted treatments for SM
- To determine the molecular epidemiology of *KIT* point mutations such as D816V, D816Y, D816F, D816H and D816L, as well as co-mutation of *KIT* with other genes such as *TET2*, *SRSF2*, *ASXL1* and *RUNX1* to investigate the molecular profiles of potential SM cases

## METHODS

- The SOPHiA DDM™ Platform (SOPHiA GENETICS SA, Switzerland) proprietary algorithms were used to analyze pseudo-anonymized real-world genomic profiles (Q1 2019 - Q2 2021) across 59 Institutions
- Data were obtained from testing with 34 SOPHiA GENETICS somatic onco-hematological (HemOnc) NGS panels capable of detecting *KIT* alterations from RNA or DNA



## RESULTS

### *KIT* testing landscape

- 32,052 individuals were tested with somatic HemOnc panels capable of detecting *KIT* alterations
- For 76% of individuals, DNA/RNA was extracted from peripheral blood samples, and for 23% of individuals from formalin-fixed paraffin-embedded (FFPE) samples
- The HemOnc somatic *KIT* testing footprint increased over time (Figure 1)
- Most cancer institutes used custom panels, while general hospitals (the majority of institutions) used more commercially available panels

### *KIT* D816 mutation disease associations

- KIT* D816 mutations were associated with a myeloid/myeloproliferative malignancy in 85% of cases with a disease tag
- Mastocytosis was the disease tag in 2% of individuals with a *KIT* D816 mutation detected by a HemOnc panel (Figure 2)
- It is important to note that the most common form of advanced SM is SM with associated hematologic neoplasm (SM-AHN), which could have been mistakenly classified under other tags such as myeloid neoplasm or leukemia




Figure 1. Number of individuals profiled with HemOnc panels covering *KIT* mutations, by country and quarter year<sup>1</sup>

<sup>1</sup>Results are extracted from SOPHiA's community and thus data should be interpreted with caution as they might reflect only part of the general trend in those countries




Figure 2. Disease tag associated with *KIT* D816 mutation-positive individuals, separated according to specific *KIT* variant (n = 283).

### *KIT* variant breakdown

- A total of 33,289 *KIT* gene variants were identified amongst 12,840 *KIT* mutation-positive individuals
- KIT* D816 variants represented 1.6-3.3% of all *KIT* mutations (Figure 3)
- KIT* D816V represented 91% of detected *KIT* D816 variants
- Seven *KIT* D816V mutation-positive cases showed intratumoral heterogeneity for *KIT* D816 variants, which could reflect a subclonal architecture of tumors

### Oncogenic alterations co-occurring with *KIT* D816V

- The analysis of *KIT* drivers within *KIT* or other genes found that 48.8% of *KIT* D816V mutation-positive cases had at least one co-occurring oncogenic alteration in *KIT*, *SRSF2*, *ASXL1*, *TET2*, or *RUNX1*
- Co-occurring oncogenic alterations were most frequent in *ASXL1* (Figure 4)




Figure 3. Frequency of specific *KIT* D816 mutations detected by HemOnc panels as a percentage of individuals positive for any *KIT* mutation, by country (normalized per number of *KIT* mutation-positive individuals per country) (total n = 12,840, n = 283 with *KIT* D816 variant).

GAM Classes	ASXL1_OG [3547]	D816 [33]	D816V [258]	KIT_OG [141]	RUNX1_OG [2192]	SRSF2_OG [2438]	TET2_OG [851]
TET2_OG [851]	112	0	8	5	52	175	621
SRSF2_OG [2438]	736	1	49	13	396	2412	175
RUNX1_OG [2192]	430	2	34	12	2048	396	52
KIT_OG [141]	22	4	9	136	12	13	5
D816V [258]	62	7	258	9	34	49	8
D816 [33]	3	32	7	4	2	1	0
ASXL1_OG [3547]	3374	3	62	22	430	736	112

Figure 4. Number of cases with oncogenic variants co-occurring with *KIT* D816V.

## CONCLUSIONS

- The comprehensive characterization of the molecular epidemiology of *KIT* variants and co-mutations is crucial to better define SM prognosis and treatment strategies
- This study provides new insights into the occurrence of *KIT* alterations and the concurrent presence of oncogenic co-mutations in potential SM cases, and how specific somatic NGS applications, namely HemOnc panels, are used across France, Italy, Germany, and Spain
- Indeed, the reported frequency for *KIT* and co-occurring alterations is likely under-estimated here, due to the availability of alternative genetic testing techniques and platforms (e.g. ddPCR, which can have a higher sensitivity than NGS)
- The low percentage of SM disease tags associated with *KIT* D816 cases highlights the difficulty in detecting and diagnosing individuals with SM and the potential number of individuals that could benefit from *KIT*-targeted treatment strategies

## REFERENCES

- Garcia-Montero AC, et al. Blood 2006;108(7):2366-72.
- Verstovsek S. Eur J Haematol. 2013;90(2):89-98.
- Gilreath JA, et al. Clin Pharmacol. 2019;11:77-92.
- Jawhar M, et al. Leukemia 2015;29(5):1115-22.
- Schwaab J, Blood 2013;122(14):2460-6.

## ACKNOWLEDGEMENTS

Scientific writing and editorial support was provided by Mallory Gough, SOPHiA GENETICS Limited.  
 Legal entity responsible for the study: SOPHiA GENETICS SA.  
 Funding: Blueprint Medicines Corporation.

## CONTACT INFORMATION

Blueprint Medicines: [medinfo@blueprintmedicines.com](mailto:medinfo@blueprintmedicines.com)  
 SOPHiA GENETICS: Sofia Lima, [SRMLima@sophiagenetics.com](mailto:SRMLima@sophiagenetics.com)